

A NOTE ON AN ASYMPTOTICALLY GOOD TAME TOWER

SIMAN YANG

ABSTRACT. The explicit construction of function fields tower with many rational points relative to the genus in the tower play a key role for the construction of asymptotically good algebraic-geometric codes. In 1997 Garcia, Stichtenoth and Thomas [6] exhibited two recursive asymptotically good Kummer towers over any non-prime field. Wulftange determined the limit of one tower in his PhD thesis [13]. In this paper we determine the limit of another tower [14].

Keywords: Function fields tower, rational places, genus.

1. INTRODUCTION

Let $K = \mathbb{F}_q$ be the finite field of cardinality q , and let $\mathcal{F} = (F_i)_{i \geq 0}$ be a sequence of algebraic function fields each defined over K . If $F_i \subsetneq F_{i+1}$ and K is the full constant field for all $i \geq 0$, and $g(F_j) > 1$ for some $j \geq 0$, we call $\mathcal{F}$ a tower.

Denoted by $g(F)$ the genus of the function field $F/\mathbb{F}_q$ and $N(F)$ the number of $\mathbb{F}_q$ -rational places of F . It is well-known that for given genus g and finite field $\mathbb{F}_q$, the number of $\mathbb{F}_q$ -rational places of a function field is upper bounded due to the Weil's theorem (cf. [11]). Let $N_q(g) := \max\{N(F) | F \text{ is a function field of genus } g \text{ over } \mathbb{F}_q\}$ and let

$$A(q) = \limsup_{g \rightarrow \infty} N_q(g)/g,$$

the Drinfeld-Vladut bound [2] provides a general upper bound of $A(q)$

$$A(q) \leq \sqrt{q} - 1.$$

Ihara[7], and Tsfasman, Vladut and Zink [12] independently showed that this bound is met when q is a square by the theory of Shimura modular curves and elliptic modular curves, respectively. For non-square q the exact value of $A(q)$ is unknown. Serre[10] first showed that $A(q)$ is positive for any prime power q

$$A(q) \geq c \cdot \log q$$

with some constant $c > 0$ irrelevant to q . It was proved in [6] that for any tower $\mathcal{F} = (F_i)_{i \geq 0}$ defined over $\mathbb{F}_q$ the sequence $N(F_n)/g(F_n)_{n \geq 0}$ is convergent. We define the limit of the tower as

$$\lambda(\mathcal{F}) = \lim_{i \rightarrow \infty} N(F_i)/g(F_i).$$

The author was partially supported by the program for Chang Jiang Scholars and Innovative Research Team in University.

Clearly, $0 \leq \lambda(\mathcal{F}) \leq A(q)$. We call a tower $\mathcal{F}$ asymptotically good if $\lambda(\mathcal{F}) > 0$. To be useful towards the aim of yielding asymptotically good codes, a tower must be asymptotically good. Practical implementation of the codes also requires explicit equations for each extension step in the tower. In 1995, Garcia and Stichtenoth [4] exhibited the first explicit tower of Artin-Schreier extensions over any finite field of square cardinality which met the upper bound of Drinfeld and Vladut. In 1997 Garcia, Stichtenoth and Thomas [6] exhibited two explicit asymptotically good Kummer towers over any non-prime field which were later generalized by Deolalikar [1]. For other explicit tame towers, readers may look at [3], [5], [9]. The two asymptotically good Kummer towers in [6] are given as below.

Let $q = p^e$ with $e > 1$, and let $F_n = \mathbb{F}_q(x_0, \dots, x_n)$ with

$$x_{i+1}^{\frac{q-1}{p-1}} + (x_i + 1)^{\frac{q-1}{p-1}} = 1 \quad (i = 0, \dots, n-1). \quad (1)$$

Then $\mathcal{F} = (F_0, F_1, \dots)$ is an asymptotically good tower over $\mathbb{F}_q$ with $\lambda(\mathcal{F}) \geq 2/(q-2)$.

Let q be a prime power larger than two, and let $F_n = \mathbb{F}_q(x_0, \dots, x_n)$ with

$$x_{i+1}^{q-1} + (x_i + 1)^{q-1} = 1 \quad (i = 0, \dots, n-1). \quad (2)$$

Then $\mathcal{F} = (F_0, F_1, \dots)$ is an asymptotically good tower over $\mathbb{F}_{q^2}$ with $\lambda(\mathcal{F}) \geq 2/(q-2)$.

Wulftange showed in [13] that $\lambda(\mathcal{F}) = 2/(q-2)$ for the first tower, we will show in the next section that the limit of the second tower is also $2/(q-2)$.

2. THE LIMIT OF THE TOWER

Lemma 2.1. *Let $F_1 = K(x, y)$ defined by Eq. (2).*

Over $K(x)$ exactly the zeroes of $x - \alpha$, $\alpha \in \mathbb{F}_q \setminus \{-1\}$ are ramified in F_1 , each of ramification index $q-1$.

Over $K(y)$ exactly the zeroes of $y - \alpha$, $\alpha \in \mathbb{F}_q^*$ are ramified in F_1 , each of ramification index $q-1$.

Proof. By applying the theory of Kummer extension (cf. [11, Chap. III.7.3]). $\square$

Proposition 2.2. *Let $P_\alpha \in \mathcal{P}(F_0)$ be a zero of $x_0 - \alpha$, $\alpha \in \mathbb{F}_q \setminus \{-1\}$. Then, P_α is totally ramified in F_{n+1}/F_n for any $n \geq 0$.*

Proof. Let $P \in \mathcal{P}(F_n)$ lying above P_α for some $\alpha \in \mathbb{F}_q \setminus \{-1\}$. From Eq. (2), one can check $x_1(P) = x_2(P) = \dots = x_n(P) = 0$. Thus the ramification index of the extension of the restriction P in $K(x_i, x_{i+1})/K(x_i)$ is $q-1$ for $i = 0, 1, \dots, n$, also the ramification index of the extension of the restriction P in $K(x_i, x_{i+1})/K(x_{i+1})$ is 1 for $i = 0, 1, \dots, n$. The proof is finished by diagram chasing and repeated application of Abhyankar's lemma. $\square$

Let $Q \in \mathcal{P}(F_n)$ be a place ramified in F_{n+1} . Then $P := Q \cap K(x_n)$ is ramified in $K(x_n, x_{n+1})$ due to Abhyankar's lemma. From Lemma 2.1, $x_n(P) = \alpha$ for some $\alpha \in \mathbb{F}_q \setminus \{-1\}$. If $\alpha \neq 0$, P is ramified in $K(x_{n-1}, x_n)$ of ramification index $q-1$ due to Lemma 2.1, and due to Abhyankar's lemma, the place in $K(x_{n-1}, x_n)$ lying above P is unramified in $K(x_{n-1}, x_n, x_{n+1})$, again by Abhyankar's lemma, Q is

unramified in F_{n+1} . Thus Q is a zero of x_n . This implies Q is a zero of $x_{n-1} - \beta$ for some $\beta \in \mathbb{F}_q \setminus \{-1\}$. From Eq. (2), one has the following possibilities for a place $Q \in \mathcal{P}(F_n)$ ramified in F_{n+1} .

- (a) The place Q is a common zero of $x_0, x_1, \dots, x_n$.
 - (b) There is some t , $-1 \leq t < n-1$ such that
 - (b1) Q is a common zero of $x_{t+2}, x_{t+3}, \dots, x_n$.
 - (b2) Q is a zero of $x_{t+1} - \alpha$ for some $\alpha \in \mathbb{F}_q^* \setminus \{-1\}$.
 - (b3) Q is a common zero of $x_0 + 1, x_1 + 1, \dots, x_t + 1$.
- (Note that condition (b2) implies (b1) and (b3)).

Lemma 2.3. *Let $-1 \leq t < n$ and $Q \in \mathcal{P}(F_n)$ be a place which is a zero of $x_{t+1} - \alpha$ for some $\alpha \in \mathbb{F}_q^* \setminus \{-1\}$. Then one has*

- (i) *If $n < 2t + 2$, then Q is unramified in F_{n+1} .*
- (ii) *If $n \geq 2t + 2$, then Q is ramified in F_{n+1} of ramification index $q - 1$.*

Proof. The assertion in (i) and (ii) follow by diagram chasing with the help of Lemma 2.1 and repeated applications of Abhyankar's lemma. $\square$

For $0 \leq t < \lfloor n/2 \rfloor$ and $\alpha \in \mathbb{F}_q^* \setminus \{-1\}$, set

$X_{t,\alpha} := \{Q \in \mathcal{P}(F_n) \mid Q \text{ is a zero of } x_{t+1} - \alpha\}$ and $A_{t,\alpha} := \sum_{Q \in X_{t,\alpha}} Q$. Denote

by Q_{t+1} the restriction of Q to $K(x_{t+1})$, we have $[F_n : K(x_{t+1})] = (q-1)^n$ and $e(Q|Q_{t+1}) = (q-1)^{n-t-1}$. Then $\deg A_{t,\alpha} = (q-1)^{t+1}$ follows from the fundamental equality $\sum e_i f_i = n$. Combining the above results one obtains

$$\deg \text{Diff}(F_{n+1}/F_n) = (q-1)(q-2) + \sum_{\alpha \in \mathbb{F}_q^* \setminus \{-1\}} \sum_{t=0}^{\lfloor n/2 \rfloor - 1} (q-2)(q-1)^{t+1} \quad (3)$$

$$= (q-2)(q-1)^{\lfloor n/2 \rfloor + 1}. \quad (4)$$

Now we can easily determine the genus of F_n by applying the transitivity of different exponents and Hurwitz genus formula. The result is:

$$g(F_{n+1}) = \begin{cases} (q-2)(q-1)^{n+1}/2 - (q-1)^{n/2+1} + 1, & \text{if } n \text{ is even,} \\ (q-2)(q-1)^{n+1}/2 - q(q-1)^{(n+1)/2}/2 + 1, & \text{if } n \text{ is odd.} \end{cases}$$

Thus $\gamma(\mathcal{F}) := \lim_{n \rightarrow \infty} g(F_n)/[F_n : F_0] = (q-2)/2$.

Remark 2.4. Note that from the proof of [6, Theorem 2.1 and Example 2.4], $\gamma(\mathcal{F})$ is upper bounded by $(q-2)/2$.

Next we consider the rational places in each function field F_n . First we consider places over P_∞ . It is easy to see that P_∞ splits completely in the tower. From Prop. 2.2, there's a unique $\mathbb{F}_q$ -rational place in F_n over P_α for any $\alpha \in \mathbb{F}_q \setminus \{-1\}$. Then we consider the K -rational place over P_{-1} in F_n . Let $0 \leq t < n$ and $Q \in \mathcal{P}(F_n)$ be a place which is a zero of $x_{t+1} - \alpha$ for some $\alpha \in \mathbb{F}_q^* \setminus \{-1\}$. We study the condition for such place Q to be K -rational.

Lemma 2.5. *Let Q' be a place of F_2 and Q' is a zero of $x_1 - \beta$ for some $\beta \in \mathbb{F}_q^* \setminus \{-1\}$. Then, if $\text{char}(\mathcal{F}) \neq 2$, Q' is not a $\mathbb{F}_q$ -rational place and Q' is a $\mathbb{F}_{q^2}$ -rational place if and only if $\beta = -1/2$; if $\text{char}(\mathcal{F}) = 2$, Q' is not a $\mathbb{F}_{q^2}$ -rational place.*

Proof. Note that x_2 and $x_0 + 1$ both are Q' -prime elements. Eq. (2) implies $(\frac{x_2}{x_0+1})^{q-1} = \frac{x_1}{1+x_1}$, which is equivalent to $\beta/(1+\beta)$. $(\frac{x_2}{x_0+1})^{q-1}(Q') \neq 1$ implies Q' is not $\mathbb{F}_q$ -rational, and $(\frac{x_2}{x_0+1})^{q^2-1}(Q') = 1$ if and only if $\beta = -1/2$ as $\beta \in \mathbb{F}_q^* \setminus \{-1\}$. $\square$

We generalize this result to the following proposition.

Proposition 2.6. *Assume $\text{char}(\mathcal{F})$ is odd. Fix positive integers $t \leq m$. There are $2^{t-1}(q-1)$ many $\mathbb{F}_{q^2} \setminus \mathbb{F}_q$ -rational places Q in $\mathbb{F}_{q^2}(x_{m-t}, x_{m-t+1}, \dots, x_{m+t})$ which are zeroes of $x_m - \beta$ for some $\beta \in \mathbb{F}_q^* \setminus \{-1\}$ if $q \equiv -1 \pmod{2^t}$, with each of them corresponds to a tuple $(\alpha_1, \alpha_2, \dots, \alpha_t)$ satisfying*

$$\left\{ \begin{array}{llll} x_m \equiv -1/2, & & & \\ x_{m+1}/(x_{m-1} + 1) & \equiv & \alpha_1, \text{ with } \alpha_1^2 & = -1, \\ x_{m+2}/(x_{m-2} + 1) & \equiv & \alpha_2, \text{ with } \alpha_2^2 & = -1/\alpha_1, \\ \dots & \equiv & \dots, \dots & = \dots, \\ x_{m+t-1}/(x_{m-t+1} + 1) & \equiv & \alpha_{t-1}, \text{ with } \alpha_{t-1}^2 & = -1/\alpha_{t-2}, \\ x_{m+t}/(x_{m-t} + 1) & \equiv & \alpha_t, \text{ with } \alpha_t^{q-1} & = -\alpha_{t-1}. \end{array} \right.$$

Proof. Prove by induction on t . For $t = 1$, this is the case in Lemma 2.5, here we take $\alpha_0 = 1$. For $t \geq 1$, it is easily checked $(\frac{x_{m+t+1}}{x_{m-t-1}+1})^{q-1} \equiv \frac{-x_{m+t}}{x_{m-t+1}}$ from definition. Thus, $\alpha_{t+1} \in \mathbb{F}_{q^2}$ if and only if $\alpha_t^{q+1} = 1$. By induction hypothesis on t , $\alpha_t^{q-1} = -\alpha_{t-1}$. Therefore Q is a $\mathbb{F}_{q^2}$ -rational place implies $\alpha_t^2 = -1/\alpha_{t-1}$. Note $\alpha_{t-1}^{2^{t-1}} = -1$. Let $q = 2^t k - 1$, we have $(-1)^k = 1$, thus k is even, i.e., $q \equiv -1 \pmod{2^{t+1}}$. This finishes the induction on $t + 1$. $\square$

Using this proposition and Lemma 2.3, we yield the following result.

Proposition 2.7. *Assume $\text{char}(\mathcal{F})$ is odd. Suppose $2^l \parallel (q+1)$. The number of $\mathbb{F}_{q^2}$ -rational place in F_n which is a zero of $x_m - \alpha$ ($0 < m \leq n$) for any $\alpha \in \mathbb{F}_q^* \setminus \{-1\}$ is counted as below.*

$$\left\{ \begin{array}{lll} 2^{m-1}(q-1), & \text{when} & 1 \leq m \leq n/2 \text{ and } m \leq l, \\ 0, & \text{when} & 1 \leq m \leq n/2 \text{ and } m > l, \\ 2^{n-m-1}(q-1), & \text{when} & n > m > n/2 \text{ and } n-m \leq l, \\ 0, & \text{when} & n > m > n/2 \text{ and } n-m > l, \\ q-2, & \text{when} & m = n. \end{array} \right.$$

Proof. Let $0 < m < n$ and $a = \min\{m, n-m\}$. If $a = m$ (resp. $a = n-m$), from Lemma 2.5, there exists $\mathbb{F}_{q^2}$ -rational place in F_{2m} (resp. $K(x_{2m-n}, \dots, x_n)$) with $x_m \equiv \alpha$ for some $\alpha \in \mathbb{F}_q^* \setminus \{-1\}$ if and only if $2^a \parallel (q+1)$, the number of such places is $2^{a-1}(q-1)$, and all these places totally ramified in F_n according to Lemma 2.1. $\square$

Hence, the number of $\mathbb{F}_{q^2}$ -rational place in F_n lying above P_{-1} is

$$\begin{cases} (q-1)(2^{l+1}-1), & \text{if } n > 2l, \\ (q-1)(2^{(n+1)/2}-1), & \text{if } n \text{ is odd and } n \leq 2l, \\ (q-1)(3 \times 2^{n/2-1}-1), & \text{if } n \text{ is even and } n \leq 2l. \end{cases}$$

Remark 2.8. If $\text{char}(\mathcal{F}) > 2$, among all $\mathbb{F}_{q^2}$ -rational place in F_n lying above P_{-1} , exactly $q-1$ are $\mathbb{F}_q$ -rational, corresponding to $x_n \equiv \alpha$ for some $\alpha \in \mathbb{F}_q^*$, respectively. If $\text{char}(\mathcal{F}) = 2$, from Lemma 2.5, there are exactly $q-1$ $\mathbb{F}_{q^2}$ -rational places in F_n lying above P_{-1} , which are all $\mathbb{F}_q$ -rational, corresponding to $x_n \equiv \alpha$ for some $\alpha \in \mathbb{F}_q^*$, respectively.

Next we determine the $\mathbb{F}_{q^2}$ -rational place Q in F_n lying above P_α for some $\alpha \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$. Direct calculation gives $x_1(Q) = \alpha_1$ for some $\alpha_1 \notin \mathbb{F}_q$. Similarly, $x_2(Q) = \alpha - 2, \dots, x_n(Q) = \alpha_n$, with $\alpha_i \in \overline{\mathbb{F}_q} \setminus \mathbb{F}_q$. We observe that Q is $\mathbb{F}_{q^2}$ -rational in F_n if and only if $\alpha, \alpha_1, \dots, \alpha_n$ are all in $\mathbb{F}_{q^2}$. To verify it, assume $\alpha, \alpha_1, \dots, \alpha_n$ are all in $\mathbb{F}_{q^2}$. Then Q is completely splitting in each extension F_i/F_{i-1} ($i = 1, 2, \dots, n$), with $x_i \equiv c\alpha_i$ for some $c \in \mathbb{F}_q^*$ in each place respectively.

We have $\alpha_1 \in \mathbb{F}_{q^2}$ if and only if $(1+\alpha)^{q-1} + (1+\alpha)^{1-q} = 1$. Similarly, α_i ($i = 1, 2, \dots, n-1$) $\in \mathbb{F}_{q^2}$ if and only if $(1+\alpha_{i-1})^{q-1} + (1+\alpha_{i-1})^{1-q} = 1$. Thus, Q is $\mathbb{F}_{q^2}$ -rational implies $(1+\alpha)^{q-1}, (1+\alpha_1)^{q-1}, \dots, (1+\alpha_{n-1})^{q-1}$ all are the root of $x^2 - x + 1 = 0$.

Claim. $(1+\alpha)^{q-1}, (1+\alpha_1)^{q-1}, \dots, (1+\alpha_{n-1})^{q-1}$ are equal.

Proof of the claim. Prove by contradiction. For simplicity assume $(1+\alpha)^{q-1} \neq (1+\alpha_1)^{q-1}$. Thus $x^2 - x + 1 = (x - (1+\alpha)^{q-1})(x - (1+\alpha_1)^{q-1})$. Comparing the coefficient of x^1 , one has $1 = (1+\alpha)^{q-1} + (1+\alpha_1)^{q-1} = (2+\alpha_1 - \alpha_1^{q-1})/(1+\alpha_1)$. This implies $\alpha_1 \in \mathbb{F}_q$, which is a contradiction.

Let $p = \text{char}(\mathbb{F}_q)$, we consider the following two cases respectively.

Case 1: $p = 3$.

Since the unique root of $x^2 - x + 1 = 0$ is -1 , $1 - \alpha_1^{q-1} = -1$, and $(1+\alpha_1)^{q-1} = -1$. It is easily checked these two equalities lead to a contradiction.

Case 2: $p \neq 3$.

Thus, -1 is not a root of $x^2 - x + 1 = 0$, which implies $(1+\alpha)^{q-1}$ and $(1+\alpha)^{1-q}$ are distinct roots of $x^2 - x + 1 = 0$. Thus, $(1+\alpha)^{q-1} + (1+\alpha)^{1-q} = 1$. By assuming $(1+\alpha)^{q-1} + (1+\alpha)^{1-q} = 1$, we have $x_1(Q) = \alpha_1$, with $\alpha_1^{q-1} = (1+\alpha)^{1-q}$. Hence, $\alpha_1 = c/(1+\alpha)$ for some $c \in \mathbb{F}_q^*$. Since $(1+\alpha)^{q-1} = (1+\alpha_1)^{q-1}$, direct calculation gives $c = \frac{(1+\alpha)^{2q-1} - (1+\alpha)^q}{1 - (1+\alpha)^{2q-2}}$. Iterating this procedure, we have a $\mathbb{F}_{q^2}$ -rational place Q in F_n lying above P_α for some $\alpha \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ is one-one corresponding to a tuple

$(c_1, c_2, \dots, c_n)$ satisfying

$$\left\{ \begin{array}{ll} x_0 \equiv \alpha, & \text{with } (1 + \alpha)^{q-1} + (1 + \alpha)^{1-q} = 1, \\ x_1 \equiv c_1/(1 + \alpha) := \alpha_1 & \text{with } c_1 = \frac{(1+\alpha)^{2q-1} - (1+\alpha)^q}{1 - (1+\alpha)^{2q-2}} \in \mathbb{F}_q^*, \\ x_2 \equiv c_2/(1 + \alpha_1) := \alpha_2 & \text{with } c_2 = \frac{(1+\alpha_1)^{2q-1} - (1+\alpha_1)^q}{1 - (1+\alpha_1)^{2q-2}} \in \mathbb{F}_q^*, \\ \dots & \text{with } \dots \\ x_{n-1} \equiv c_{n-1}/(1 + \alpha_{n-2}) := \alpha_{n-1} & \text{with } c_{n-1} = \frac{(1+\alpha_{n-2})^{2q-1} - (1+\alpha_{n-2})^q}{1 - (1+\alpha_{n-2})^{2q-2}} \in \mathbb{F}_q^*, \\ x_n \equiv c_n/(1 + \alpha_{n-1}), & \text{with } c_n \in \mathbb{F}_q^*. \end{array} \right.$$

Therefore, for any $\alpha \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$, the number of $\mathbb{F}_{q^2}$ -rational places in F_n lying above P_α is zero if $\text{char}(\mathcal{F}) = 3$; and $(q-1)\#\{\alpha \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q : (1+\alpha)^{q-1} + (1+\alpha)^{1-q} = 1\}$ if $\text{char}(\mathcal{F}) \neq 3$.

As we have determined all $\mathbb{F}_q$ -rational places and $\mathbb{F}_{q^2}$ -rational places in F_n , we are now able to determine the value of $\nu(\mathcal{F})$. If $\text{char}(\mathcal{F}) \neq 2$ and the constant field is $\mathbb{F}_q$, then $\nu(\mathcal{F}) = 0$, and $\nu(\mathcal{F}) = 1$ if the constant field is $\mathbb{F}_{q^2}$. If $\text{char}(\mathcal{F}) = 2$, and the constant field is $\mathbb{F}_q$ ($q > 2$), then $\nu(\mathcal{F}) = 1$.

Remark 2.9. One can check that the function field tower recursively defined by Eq. (2) is isomorphic in some extension field of $\mathbb{F}_{q^2}$, to a tower recursively defined by $y^{q-1} = 1 - (x + \alpha)^{q-1}$, where α is any nonzero element of $\mathbb{F}_q$.

From above discussion, Eq.(2) defines an asymptotically bad tower over any prime field (it does not define a tower over $\mathbb{F}_2$). Lenstra showed in [8, Theorem 2] that there does not exist a tower of function fields $\mathcal{F} = (F_0, F_1, \dots)$ over a prime field which is recursively defined by $y^m = f(x)$, where $f(x)$ is a polynomial $f(x)$, m and q are coprime, such that the infinity place of F_0 splits completely in the tower, and the set $V(\mathcal{F}) = \{P \in \mathcal{P}(F_0) | P \text{ is ramified in } F_n/F_0 \text{ for some } n \geq 1\}$ is finite. A tower recursively defined by Eq. (2) falls in this form with a finite set $V(\mathcal{F})$, but no place of F_0 splits completely in the tower. Thus arises a problem: can one find an asymptotically good, recursive tower of the above form, over a prime field, with a finite set $V(\mathcal{F})$ and a finite place splitting completely in the tower?

REFERENCES

- [1] V. Deolalikar, On splitting places of degree one in extensions of algebraic function fields, towers of function fields meeting asymptotic bounds, and basis constructions for algebraic-geometric codes. PhD thesis, University of Southern California, Los Angeles, May 1999.
- [2] V.G. Drinfeld, S.G. Vlădut, Number of points of an algebraic curve, *Func. Anal.*, **17** (1983), 53-54.
- [3] N. D. Elkies, Explicit modular towers, *Proceedings of the Thirty-Fifth [1997] Annual Allerton Conference on Communication, Control and Computing*, (T. Basar and A. Vardy, eds.), Univ. of Illinois at Urbana-Champaign, (1998), 23-32.
- [4] A. Garcia, H. Stichtenoth, A tower of Artin-Schreier extensions of function fields attaining the Drinfeld-Vladut bound, *Invent. Math.* **121** (1995), 211-222.
- [5] A. Garcia, H. Stichtenoth, On tame towers over finite fields, *J. Reine Angew. Math* **557** (2003), 53-80.
- [6] A. Garcia, H. Stichtenoth, M. Thomas, On towers and composita of towers of function fields over finite fields, *Finite fields and their applications* (3) **3** (1997), 257-274.

- [7] Y. Ihara, Some remark on the number of rational points of algebraic curves over finite fields, *J. Fac. Sci. Tokyo*, **28** (1981), 721-724.
- [8] H. W. Lenstra, Jr., On a problem of Garcia, Stichtenoth, and Thomas, *Finite fields and their applications* (2) **8** (2002), 166-170.
- [9] W.-C. W. Li, H. Maharaj, H. Stichtenoth, New optimal towers over finite fields, In: *Algorithmic Number Theory [Sydney, 2002]* (C. Fieker and D. Kohel, eds.), 372-389.
- [10] J.-P. Serre, “Rational points on curves over finite fields”, *Lecture Notes*, Harvard University, 1985.
- [11] H. Stichtenoth, “Algebraic Function Fields and Codes”, *Springer Universitext*, Springer, Berlin, 1993.
- [12] M. A. Tsfasman, S.G. Vlăduț, T. Zink, Modular curves, Shimura curves, and Goppa codes, better than the Varshamov-Gilbert bound, *Math. Nachr.* **109** (1982), 21-28.
- [13] J. Wulftange, *Türme algebraischer Funktionenkörper*, PhD thesis, Essen, 2002.
- [14] S. Yang, *On explicit towers of function fields over finite fields*, PhD thesis, National University of Singapore, 2005.

Siman Yang

Department of Mathematics, East China Normal University,
500, Dongchuan Rd., Shanghai, P.R.China 200241. e-mail: smyang@math.ecnu.edu.cn